

Cosmic Censorship in Higher Dimensions

Rituparno Goswami* and Pankaj S Joshi†

*Department of Astronomy and Astrophysics
Tata Institute of Fundamental Research
Homi Bhabha Road, Mumbai 400 005, India*

We show that the naked singularities arising in dust collapse from smooth initial data (which include those discovered by Eardley and Smarr, Christodoulou, and Newman) are removed when we make transition to higher dimensional spacetimes. The cosmic censorship is then restored for dust collapse which will always produce a black hole as the collapse end state for dimensions $D \geq 6$, under conditions to be motivated physically such as the smoothness of initial data from which the collapse develops.

PACS numbers: 04.20.Dw, 04.20.Cv, 04.70.Bw

A considerable debate has continued in recent years on the validity or otherwise of the cosmic censorship conjecture (CCC) introduced by Penrose and Hawking, which effectively states that any singularities that arise from gravitational collapse from a regular initial data must not be visible to far away observers in the spacetime and are always hidden within black holes. Such an assumption has been used extensively and is fundamental to the theory as well as applications of black hole physics. On the other hand, if a naked singularity results as collapse end state, it is no longer necessarily covered by the event horizon and could communicate, in principle, with outside observers. Such a scenario is of physical interest because a naked singularity may have theoretical and observational properties quite different from a black hole end state, and communications from extreme strong gravity regions dominated by quantum gravity may be possible.

Though there is no satisfactory proof or mathematical formulation of CCC available despite many efforts, there are many examples of dynamical collapse models available which lead to a black hole (BH) or a naked singularity (NS) as the collapse end state, depending on the nature of the initial data (see e.g. [1] and references therein). In particular, pioneering analytic models by Christodoulou, and Newman, and numerical work by Eardley and Smarr [2], established the existence of *shell-focusing* naked singularities as the end state of a continual collapse, where the physical radius of all collapsing shells vanishes. In these models the matter form was taken to be marginally bound dust, assuming that the initial data functions are smooth and even profiles. Newman generalized these models for non-marginally bound class, thus covering the entire class of dust collapse solutions. A general treatment for dust collapse with generic initial data was developed in [3].

In fact, gravitational collapse of a dust cloud has served, over past several decades, as a basic and fundamental paradigm in black hole physics. This was first

discussed by Oppenheimer and Snyder [4], who considered a continual collapse of a homogeneous spherical dust cloud in general relativity. The important feature that emerges is that as the collapse progresses, an event horizon of gravity forms, which fully covers the spacetime singularity developing later as the collapse end state. Since the singularity is always covered by the horizon, we say a black hole has developed as the collapse final state. This scenario was the basic motivation and origin for CCC, stating that any physically realistic collapse must have a similar outcome producing a black hole only.

Our purpose here is to show explicitly, that the above mentioned naked singularities of dust collapse (for both marginally bound and non-marginally bound dust) are removed when we go to higher (≥ 6) dimensions, if we allow for only smooth and analytic initial profiles as assumed by [2]. In other words, the collapse of a dust-like matter, whose initial density and velocity profiles are taken to be smooth C^∞ functions, which is a condition to be motivated physically, always goes to a black hole end state for dimensions $D \geq 6$, and CCC is restored. We point out that the increase in dimensions deforms the apparent horizon, which is the outer boundary of trapped surfaces, in such a manner that the formation of the trapped surfaces is advanced. Hence, there exists a critical dimension beyond which the neighborhood of the center gets trapped before the central singularity and the final outcome of collapse is a black hole necessarily.

Such a scenario offers a possibility to recover CCC, because it is quite possible that we may be actually living in a universe with higher dimensions. The recent developments in string theory and other field theories strongly indicate that gravity is possibly a higher dimensional interaction, which reduces to general relativistic description at lower energies. Hence, it could be that CCC is restored due to extra physical effects arising from our transition itself to a higher-dimensional spacetime continuum. The recent revival of interest in this problem is motivated partly by the Randall-Sundrum brane-world models [5].

To study the collapse of a spherical dust cloud, we choose a general spherically symmetric comoving metric

*Electronic address: goswami@tifr.res.in

†Electronic address: psj@tifr.res.in

in $N \geq 4$ dimensions which has the form,

$$ds^2 = -e^{\nu(t,r)} dt^2 + e^{2\psi(t,r)} dr^2 + R^2(t,r) d\Omega_{N-2}^2 \quad (1)$$

where,

$$d\Omega_{N-2}^2 = \sum_{i=1}^{N-2} \left[\prod_{j=1}^{i-1} \sin^2(\theta^j) \right] (d\theta^i)^2 \quad (2)$$

is the metric on $(N-2)$ sphere. The energy-momentum tensor of the dust has the form,

$$T_t^t = \rho(t,r); \quad T_r^r = 0; \quad T_{\theta^i}^{\theta^i} = 0 \quad (3)$$

We take the matter field to satisfy the *weak energy condition*, that is, the energy density measured by any local observer be non-negative, and so for any timelike vector V^i , we must have,

$$T_{ik} V^i V^k \geq 0 \implies \rho \geq 0 \quad (4)$$

In the case of a finite collapsing cloud, there is a finite boundary $0 < r < r_b$, outside which it is matched to a Schwarzschild exterior. The range of the coordinates for the metric is then $0 < r < \infty$, and $-\infty < t < t_s(r)$ where $t_s(r)$ corresponds to the singular epoch $R = 0$. Solving the N -dimensional Einstein equations we get the generalized Tolman-Bondi-Lemaître (TBL) metric [6] as,

$$ds^2 = -dt^2 + \frac{R'^2}{1+f(r)} dr^2 + R^2(t,r) d\Omega_{N-2}^2 \quad (5)$$

where $f(r)$ is an arbitrary function of the comoving radius r , and $f(r) > -1$. Also we get the required equations of motion as,

$$\frac{(N-2)F'}{2R^{(N-2)}R'} = \rho \quad ; \quad \dot{R}^2 = \frac{F(r)}{R^{(N-3)}} + f(r) \quad (6)$$

Here $F(r)$ is another arbitrary function of the comoving coordinate r . In spherically symmetric spacetimes $F(r)$ has the interpretation of *mass function* which describes the mass distribution of the dust cloud and $f(r)$ is the *energy function*, which specifies the velocity profiles for the collapsing shells for any given mass function. The energy condition then implies $F' \geq 0$. It follows from the above that there is a spacetime singularity at $R = 0$ and at $R' = 0$. The latter are called ‘shell-crossing’ singularities, which occur when successive shells of matter cross each other. These have not been considered generally to be genuine spacetime singularities, and possible extensions of spacetime have been investigated through the same [7]. On the other hand, the singularity at $R = 0$ is where all matter shells collapse to a zero physical radius, and hence this has been known as a ‘shell-focusing’ singularity. The nature of this singularity has been investigated extensively in four-dimensional spacetimes (see e.g. references in [1]), and it is known for the case of spherical dust collapse that this can be both naked or covered

depending on the nature of the initial data from which the collapse develops. The collapse condition is $\dot{R} < 0$.

We now use the scaling independence of the comoving coordinate r to write (see e.g. [8]),

$$R(t,r) = rv(t,r) \quad (7)$$

where,

$$v(t_i, r) = 1 \quad ; \quad v(t_s(r), r) = 0 \quad ; \quad \dot{v} < 0 \quad (8)$$

This means we have scaled the coordinate r in such a way that at the initial epoch $R = r$, and at the singularity $R = 0$. It should be noted that we have $R = 0$ both at the regular center $r = 0$ of the cloud, and at the spacetime singularity, where all matter shells collapse to a zero physical radius. The regular center is then distinguished from the singularity by a suitable behaviour of the mass function $F(r)$ so that the density remains finite and regular there at all times till the singular epoch. The introduction of the parameter v as above then allows us to distinguish the spacetime singularity from the regular center, with $v = 1$ at the initial epoch, including the center $r = 0$, which then decreases monotonically with time as collapse progresses to value $v = 0$ at the singularity $R = 0$.

In order to ensure the regularity of the initial data, it is evident from the equations of motion that at the initial epoch the two free functions $F(r)$ and $f(r)$ must have the following forms,

$$F(r) = r^{(N-1)} \mathcal{M}(r); \quad f(r) = r^2 b(r) \quad (9)$$

Following [2], we now assume that the initial density and energy functions $\rho(r)$ and $f(r)$ are smooth and even, ensuring their analytic nature. We note that the Einstein equations as such do not impose any such restriction, which has to be physically motivated, and it implies a certain mathematical simplicity in arguments to deal with a dynamical collapse. It follows that both $\mathcal{M}(r)$ and $b(r)$ are now smooth C^∞ functions, which means the Taylor expansions of these functions around the center must be of the following form,

$$\mathcal{M}(r) = \mathcal{M}_0 + r^2 \mathcal{M}_2 + r^4 \mathcal{M}_4 + \dots \quad (10)$$

and

$$b(r) = b_0 + r^2 b_2 + r^4 b_4 + \dots \quad (11)$$

that is, all odd terms in r vanish in these expansions, and the presence of only even terms would ensure smoothness.

To predict the final state of collapse for a given initial mass and velocity distribution, we shall study below the singularity curve resulting from the collapse of successive matter shells, and the apparent horizon developing in the spacetime. A decreasing apparent horizon in (t, r) plane then is a sufficient condition for a BH as it shows the entrapment of the neighborhood of the center before the central singularity.

Consider now the endless collapse of a dust cloud to a final shell-focusing singularity at $R = 0$, where all matter shells collapse to a zero physical radius. With the regular initial conditions as above, equation(6) can be written as,

$$v^{\frac{N-3}{2}} \dot{v} = -\sqrt{\mathcal{M}(r) + v^{(N-3)}b(r)} \quad (12)$$

Here the negative sign implies that $\dot{v} < 0$, that is, the matter cloud is collapsing. Integrating the above equation with respect to v , we get,

$$t(v, r) = \int_v^1 \frac{v^{\frac{N-3}{2}} dv}{\sqrt{\mathcal{M}(r) + v^{(N-3)}b(r)}} \quad (13)$$

We note that the coordinate r is to be treated as a constant in the above equation. Expanding $t(v, r)$ around the center, we get,

$$t(v, r) = t(v, 0) + r\mathcal{X}(v) + r^2 \frac{\mathcal{X}_2(v)}{2} + r^3 \frac{\mathcal{X}_3(v)}{6} + \dots \quad (14)$$

Now from equation(10) and (11) we have $\mathcal{X}(v) = 0$. The function $\mathcal{X}_2(v)$ is then given as,

$$\mathcal{X}_2(v) = - \int_v^1 \frac{v^{\frac{N-3}{2}} (\mathcal{M}_2 + v^{(N-3)}b_2) dv}{(\mathcal{M}_0 + v^{(N-3)}b_0)^{\frac{3}{2}}} \quad (15)$$

where,

$$b_0 = b(0); \mathcal{M}_0 = \mathcal{M}(0); b_2 = b''(0); \mathcal{M}_2 = \mathcal{M}''(0) \quad (16)$$

We note that the value of $\mathcal{X}_2$ is determined fully by the initial values of the mass function $\mathcal{M}(r)$ and the energy function $b(r)$. Thus, the time taken for the central shell to reach the singularity is given by

$$t_{s_0} = \int_0^1 \frac{v^{\frac{N-3}{2}} dv}{\sqrt{\mathcal{M}_0 + v^{(N-3)}b_0}} \quad (17)$$

From the above equation it is clear that for t_{s_0} to be defined,

$$\mathcal{M}_0 + v^{(N-3)}b_0 > 0 \quad (18)$$

that is, the continual collapse condition implies the positivity of the above term. Hence the time taken for other shells close to the center to reach the singularity can now be given by,

$$t_s(r) = t_{s_0} + r^2 \frac{\mathcal{X}_2(0)}{2} + \dots \quad (19)$$

In order to determine the visibility or otherwise of the singularity at $R = 0$, we need to analyze the causal structure of the trapped surfaces and the nature and behaviour of null geodesics in the vicinity of the same. If there exist future directed null geodesics with past end point at the singularity, which go out to faraway observers in space-time, then the singularity is naked. In the case otherwise,

we have a black hole resulting as the end state of a continual collapse. The boundary of the trapped region of the space-time is given by the apparent horizon within the collapsing cloud, which is given by the equation,

$$\frac{F}{R^{N-3}} = 1 \quad (20)$$

Broadly, it can be stated that if the neighborhood of the center gets trapped earlier than the singularity, then it is covered, otherwise it is naked with families of non-spacelike future directed trajectories escaping away from it. For example, it follows from the above equation that along the singularity curve $t = t_s(r)$ (which corresponds to $R = 0$), for any $r > 0$ we have $F(r)$ going to a constant positive value, whereas $R \rightarrow 0$. Hence it follows that trapping already occurs before the singularity develops at any $r > 0$ along the singularity curve $t_s(r)$.

What we need to determine now is when there will be families of non-spacelike paths coming out of the central singularity at $r = 0, t = t_s(0)$, reaching outside observers, and when there will be none. The visibility or otherwise of the singularity is decided accordingly. By determining the nature of the singularity curve, and its relation to the initial data, we are able to deduce whether the trapped surface formation in collapse takes place before or after the central singularity. It is this causal structure that determines the possible emergence or otherwise of non-spacelike paths from the singularity, and settles the final outcome in terms of either a BH or NS. From equations(20), we have,

$$v_{ah}(r) = [r^2 \mathcal{M}(r)]^{\frac{1}{N-3}} \quad (21)$$

Using the above equation in (13) we get

$$t_{ah}(r) = t_s(r) - \int_0^{v_{ah}(r)} \frac{v^{\frac{N-3}{2}} dv}{\sqrt{\mathcal{M}(r) + v^{(N-3)}b(r)}} \quad (22)$$

As we are considering the behaviour of the apparent horizon close to the central singularity at $r = 0, R = 0$ (all other points $r > 0$ on the singularity curve are already covered, as we pointed out above), therefore the upper limit of integration in the above equation is small, and hence we can expand the integrand in a power series in v , and keep only the leading order term, which amounts to,

$$t_{ah}(r) = t_{s_0} + r^2 \frac{\mathcal{X}_2(0)}{2} + \dots - r^{\frac{N-1}{N-3}} \frac{2}{N-1} \mathcal{M}_0^{\frac{1}{N-3}} \quad (23)$$

Since the apparent horizon is a well-behaved surface for a spherical dust collapse, hence we can say that the singularity curve for the collapse and its derivatives around the center are also well-defined, as the same coefficients are present in both (23) and (14).

It is now possible to analyze the effect of the number of dimensions on the nature and shape of the apparent horizon. As we increase the number of dimensions and go to dimensions higher than five, the negative term




FIG. 1: The apparent horizon in different spacetime dimensions. Here $\mathcal{M}(r) = 1 - 3.6r^2 + \dots$ and $b(r) = 1 - 3.6r^2 + \dots$, and apparent horizon curves are given for dimensions 4 to 7.

in equation(23) starts dominating, thus advancing the trapped surface formation in time. Fig. 1 shows the behaviour of the apparent horizon curve for the same initial data in different dimensions. As seen here, in usual four-dimensional spacetime the above initial profile ensures an increasing apparent horizon and it can be explicitly shown that the end state is a NS. But in six and higher spacetime dimensions we see the negative term starts dominating, purely as a result of increase in dimensions. This causes the trapped surfaces to form sufficiently early to cover the singularity. It follows that the number of spacetime dimensions causes an intriguing effect on the causal structure and nature of the apparent horizon, as seen above clearly.

We thus see that for a smooth initial data and for dimensions higher than five, the apparent horizon becomes a decreasing function of r near the center. This implies that the neighborhood of the center gets trapped before the central singularity and the central singularity is then

always covered, as opposed to the four-dimensional models by Eardley and Smarr, Christodoulou and Newman, in which case a positive $\mathcal{X}_2(0)$ would make the apparent horizon curve increasing and hence ensures a naked singularity. To be specific, suppose there is a future directed outgoing null geodesic coming out from the central singularity at $R = 0, r = 0$. If (t_1, r_1) is an event along the same, then $t_1 > t_{s_0}$ and $r_1 > 0$. But for any such r_1 , the trapped region already starts before $t = t_{s_0}$, hence the event (t_1, r_1) is already in the trapped region and the geodesic cannot be outgoing. Thus, there are no outgoing paths from the central singularity, making it covered. It now follows in general (i.e. for both marginally bound and non-marginal cases) that *for a dust collapse with smooth initial profiles, the final outcome is always a black hole for any spacetime dimensions $D \geq 6$, and all the naked singularities occurring in lower dimensions are removed.*

In five dimensions we have an interesting scenario arising (see also [9]). As it is clearly seen from the equation(23), we have a critical value of $\mathcal{X}_2(0)$, below which the apparent horizon is decreasing and we will get a BH end state. However, in the case otherwise, a naked singularity can result. It is interesting to note also that the above results holds if the initial profiles, instead of being absolutely smooth C^∞ functions, are taken to be sufficiently smooth (i.e. at least a C^2 function).

We note that it may still be possible to have both BH/NS outcomes as collapse end states when initial data and metric are not assumed necessarily to be C^∞ or at least C^2 analytic functions [3]. Hence such smoothness conditions need to be investigated and probed further carefully to see if they could be strongly motivated from a physical perspective. It will also be interesting to see the effects of introducing pressures within the collapsing cloud, and to examine if we could still avoid naked singularities and preserve cosmic censorship in higher dimensions by introducing physically motivated conditions such as above. This will be discussed elsewhere.

-
- [1] A. Krolak, Prog. Theor. Phys. Suppl. **136**, 45 (1999).
R. Penrose, in *Black holes and relativistic stars*, ed. R. M. Wald (University of Chicago Press, 1998); R. M. Wald, gr-qc/9710068; P. S. Joshi, Pramana **55**, 529 (2000); M. Celerier and P. Szekeres, Phys.Rev. D65 (2002) 123516; R. Giambo', F. Giannoni, G. Magli, P. Piccione, Commun.Math.Phys. 235 (2003) 545-563; T. Harada, H. Iguchi, and K. Nakao, Prog.Theor.Phys. 107 (2002) 449-524; T. Hertog, G. T. Horowitz and K. Maeda, gr-qc/0307102; G. Horowitz, hep-th/0312123.
 - [2] D. M. Eardley and L. Smarr, Phys. Rev. D 19, p.2239 (1979); D. Christodoulou, Commun. Math. Phys., 93, p.171 (1984); R. P. A. C. Newman, Class. Quantum Grav., 3, p.527 (1986).
 - [3] P. S. Joshi and I. H. Dwivedi, Phys. Rev. D 47, p.5357 (1993).
 - [4] J. Oppenheimer and H. Snyder, Phys. Rev. 56, p.455 (1939).
 - [5] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690(1999).
 - [6] R. C. Tolman, Proc. Natl. Acad. Sci. USA, 20, p.410 (1934); H. Bondi, Mon. Not. Astron. Soc., 107, p.343 (1947); G. Lemaitre, Ann. Soc. Sci.Bruxelles I, **A53**, 51(1933).
 - [7] C. J. S. Clarke, *The analysis of spacetime singularities*, Cambridge University Press, Cambridge, 1993.
 - [8] P. S. Joshi and I. H. Dwivedi, Class. Quantum Grav. **16**, 41 (1999).
 - [9] A. Banerjee, U. Debnath, S. Chakraborty, gr-qc/0211099; K.D. Patil, S.H. Ghate, and R.V. Saraykar, Ind. Jour. of Pure and App.Maths., Vol. 33, No. 3 (2002) p. 379; S. G. Ghosh and A. Beesham, Phys.Rev. **D64**, 124005(2001).